

Contrast-enhanced multiparametric prostate resonance

REQUIREMENTS AND PREPARATION



Requirements for the procedure:



The patient must inform at the time of assigning the appointment if they have brackets, splints, stents, external tutors, valves, pacemakers, splinters or other metallic elements.



The patient or relative must inform if they suffer from claustrophobia or involuntary movements

If the MRI is contrasted, you must take into account: inform if you have a history of allergies or known



The patient must be accompanied by an adult on the day of the exam.

reactions to contrast media, diabetes and kidney failure.



If the hospitalized patient must bring a canalized vein and come with an adult relative.

(Diagnosis: Parkinson's, dementia and Alzheimer's). Some cases require anesthesia.



If you have had a prostate biopsy, the appointment should be scheduled one month after it was performed.



The patient and his companion should not bring metallic objects (chains, rings, earrings, etc.).

Instructions for preparation:



Fast for six (6) hours. The meal before fasting should not contain fat, dairy, dark drinks or carbonated drinks



Do not suspend control medications, they should be taken only with three (3) sips of water.



The patient should apply a rectal Travad enema two (2) hours prior to the procedure.

Instructions for the administration of TRAVAD products





Before starting administration, carefully read the instructions for use.


ENEMA TRAVAD x 133 ML



Remove the protector from the blue rectal cannula (a).



Lie on one side and insert the cannula (a) gently into the rectum.



Break the green valve (b) by flexing it up and down repeatedly.



Squeeze bag with one hand until contents are completely empty.



Remove the cannula (a), retain the fluid for 3 to 5 minutes, then evacuate.

Documents required for the procedure:



Identification document.



Current order (does not apply to individuals).



Complete Clinical History with data justifying the study.



Resonances, Tomographies, Ultrasounds and previous studies.



Blood creatinine test not exceeding thirty days (30 days).

Be half an hour (30 minutes) before the procedure

This space will be necessary to register your information and confirm the correct preparation for the performance of your procedure.



Important:

If for any reason it is not possible for you to attend your appointment or require additional information before or after it, please call: (604) 444 5690 option 5.

