



# Contrast-enhanced multiparametric prostate resonance

## REQUIREMENTS AND PREPARATION



### Requirements for the procedure:

|  |                                                                                                                                                                                     |
|--|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | The patient must inform at the time of assigning the appointment if they have brackets, splints, stents, external tutors, valves, pacemakers, splinters or other metallic elements. |
|  | If the MRI is contrasted, you must take into account: inform if you have a history of allergies or known reactions to contrast media, diabetes and kidney failure.                  |
|  | The patient or relative must inform if they suffer from claustrophobia or involuntary movements (Diagnosis: Parkinson's, dementia and Alzheimer's). Some cases require anesthesia.  |
|  | The patient must be accompanied by an adult on the day of the exam.                                                                                                                 |
|  | If the hospitalized patient must bring a canalized vein and come with an adult relative.                                                                                            |
|  | If you have had a prostate biopsy, the appointment should be scheduled one month after it was performed.                                                                            |
|  | The patient and his companion should not bring metallic objects (chains, rings, earrings, etc.).                                                                                    |

### Instructions for preparation:

|  |                                                                                             |
|--|---------------------------------------------------------------------------------------------|
|  | Do not suspend control medications, they should be taken only with three (3) sips of water. |
|  | The patient should apply a rectal Travad enema two (2) hours prior to the procedure.        |

### Instructions for the administration of TRAVAD products



Before starting administration, carefully read the instructions for use.

#### ENEMA TRAVAD x 133 ML

|   |   |   |   |   |
|---|---|---|---|---|
|   |   |   |   |   |
| 1 | 2 | 3 | 4 | 5 |

### Documents required for the procedure:

|  |                                                           |  |                                                             |
|--|-----------------------------------------------------------|--|-------------------------------------------------------------|
|  | Identification document.                                  |  | Resonances, Tomographies, Ultrasounds and previous studies. |
|  | Current order (does not apply to individuals).            |  | Blood creatinine test not exceeding thirty days (30 days).  |
|  | Complete Clinical History with data justifying the study. |  |                                                             |

#### Be half an hour (30 minutes) before the procedure

This space will be necessary to register your information and confirm the correct preparation for the performance of your procedure.

ATTEND ON TIME

### Important:

If for any reason it is not possible for you to attend your appointment or require additional information before or after it, please call: (604) 444 5690 option 5.